

Orient Pharm Exp Med (2011) 11:77–90
DOI 10.1007/s13596-011-0019-1

REVIEW

A review on herbal antiasthmatics

Ravindra G. Mali · Avinash S. Dhake

Received: 27 January 2011 / Accepted: 1 April 2011 / Published online: 5 July 2011
© Institute of Oriental Medicine, Kyung Hee University 2011

Abstract In traditional systems of medicine, many plants have been documented to be useful for the treatment of various respiratory disorders including asthma. In the last two decades the use of medicinal plants and natural products has been increased dramatically all over the world. Current synthetic drugs used in pharmacotherapy of asthma are unable to act at all the stages and targets of asthma. However some herbal alternatives employed in asthma are proven to provide symptomatic relief and assist in the inhibition of disease progression also. The herbs have shown interesting results in various target specific biological activities such as bronchodilation, mast cell stabilization, anti-anaphylactic, anti-inflammatory, anti-spasmodic, anti-allergic, immunomodulatory and inhibition of mediators such as leukotrienes, lipoxygenase, cyclooxygenase, platelet activating, phosphodiesterase and cytokine, in the treatment of asthma. This paper is an attempt to classify these pharmacological and clinical findings based on their possible mechanism of action reported. It also signifies the need for development of polyherbal formulations containing various herbs acting at particular sites of the patho-

physiological cascade of asthma for prophylaxis as well as for the treatment of asthma.

Keywords Asthma · Current therapy · Herbal therapy · Poly herbal formulations · Ayurvedic drugs · Medicinal plants

Introduction

According to the National Institute of Health (NIH), asthma is defined as a chronic inflammatory disorder of the airways in which many cells and cellular elements play a role, in particular, mast cells, eosinophils, T-lymphocytes, neutrophils and epithelial cells (NIH 1997). Asthma is caused by a very complex interaction between inflammatory cells and mediators. Herbal approaches have regained their popularity, for the treatment of asthma, with their efficacy and safety aspects being supported by controlled clinical studies (Huntley and Ernst 2000). Ongoing worldwide research has also provided valuable clues regarding the precise mechanism of action of these herbal alternatives (Goyal et al. 2007).

Pharmacotherapy of bronchial asthma

In the past most clinicians managed asthma mainly according to the patient's symptom. Asthma was regarded primarily as a problem of bronchospasm and measures to prevent or reverse bronchospasm comprised the mainstay of therapy. However, during early 1980s when asthma emerged as an inflammatory rather than primarily a bronchospastic disorder, the basic approach switched from control of symptoms to control of underlying airway inflammation (Barns 1989). According to guidelines of The National Asthma Education and Prevention Program's

R. G. Mali
L. B. Rao Institute of Pharmaceutical Education and Research,
Shri B. D. Rao College Campus,
Khambhat 388 620, India

A. S. Dhake
S.M.B.T. College of Pharmacy, Dhamangaon,
Tal: Igatpuri,
Nashik 422 403, India

R. G. Mali (✉)
Department of Pharmacognosy,
L. B. Rao Institute of Pharmaceutical Education and Research,
Khambhat 388 620 Gujarat, India
e-mail: ravigmali@yahoo.co.in

(NAEPP) guidelines for the diagnosis and management of asthma, the treatment should have following goals:

1. Maintain normal activity levels, including exercise.
2. Maintain normal or near normal pulmonary function.
3. Prevent chronic and troublesome symptoms.
4. Prevent recurrent exacerbations.
5. Avoid adverse effects from medications.

The pharmacological management of asthma depends upon frequency and severity of patient's symptoms. Infrequent attacks can be managed by treating each attack when it occurs, but with more frequent attacks preventive therapy needs to be used. The following categories of drugs are used in asthma:

1. Bronchodilators
 - 1.1 β -adrenergic agonists: e.g. Metaproterenol, terbutaline, albuterol, formoterol, bitolterol, salmeterol, pirbuterol.
 - 1.2 Anticholinergics: e.g. Ipratropium bromide, Tiotropium bromide.
 - 1.3 Methylxanthines: e.g. Theophylline, aminophylline, acepihylline, diprophylline, proxophylline.
2. Anti-inflammatory agents
 - 2.1 Corticosteroids: e.g. Prednisolone, dexamethasone, beclomethasone dipropionate, dexamethasone, budesonide, fluticasone.
 - 2.2 Anti-leukotrienes: e.g. Probilukast, Irelukast, Zileuton, Montelukast, zafirlukast, pranlukast.
 - 2.3 Mast Cell Stabilizers: e.g. Cromolyn Sodium, Nedocromil sodium.

Bronchodilators

Bronchodilator drugs have an anti-bronchoconstrictor effect that may be demonstrated directly in vitro by drug-induced relaxation of precontracted airways (Barns et al. 1988). Bronchodilators promptly reverse airway obstruction in asthmatics. This action believed to be mediated by a direct effect on airway smooth muscle. However, additional pharmacologic effects on the other airway cells (such as capillary endothelium to reduce microvascular leakage and mast cells to reduce release of bronchoconstrictor mediators) may contribute to the overall reduction in airway narrowing. Only three types of bronchodilators are in current clinical use: β -adrenergic agonists, methylxanthines, and anticholinergics.

β -adrenergic agonists

Epinephrine has been used to treat asthma since the beginning of the 20th century. β Adrenergic agonists are

most widely used and effective bronchodilators for the treatment of asthma. Bronchodilation is mediated by β_2 receptors; β_2 selective drugs (Salmeterol and Formoterol) have been developed that have long duration of effect. β Adrenergic agonists lead to relaxation of bronchial smooth muscle that promote bronchodilation. Activation of adenylylate cyclase increases the concentration of intracellular cyclic adenosine 3', 5'-monophosphate (cAMP), leading to activation of specific cAMP-dependent protein kinases that cause relaxation. Relaxation may also be due to inhibition of myosin phosphorylation. β -adrenergic agonists reverse bronchoconstriction irrespective of the contractile agent. β -adrenergic agonists prevent release of mediators from a number of inflammatory cells in vitro (Church and Hiroi 1987). In addition, β adrenergic agonists increase mucus secretion from submucosal glands and ion transport across airway epithelium. These effects enhance mucociliary clearance caused by asthma (Pavia et al. 1980).

The inhaled route of administration is preferable to the oral route because adverse effects caused by systemic action of the drug are less and also because this route may be more effective. The inhaled drug reaches surface cells (e.g., mast cells or epithelial cells), which are less accessible to the orally administered drug.

Metaproterenol, terbutaline, albuterol, formoterol, bitolterol, salmeterol, and pirbuterol are the classic examples of selective β_2 -adrenergic agonists.

β agonists improve respiratory symptoms and exercise tolerance despite the small improvement in spirometric measurements. The long acting β -agonists decrease infection exacerbations as an additional potential benefit. Salmeterol has been shown to reduce adherence of bacteria such as *H. influenza* to airway epithelial cells.

β_2 selective agents cause tachycardia and palpitation by reflex cardiac stimulation secondary to peripheral vasodilation. Muscle tremor is caused by stimulation of β_2 adrenergic receptors in skeletal muscle and is the primary adverse effect of albuterol and bitolterol. Transient hypokalemia may be induced by high dose of these agents.

Anticholinergics

Datura plants contain the muscarinic antagonist and were smoked for relief of asthma centuries ago. Now a days, atropine and ipratropium bromide are the most commonly available anticholinergics.

Antimuscarinic agents specifically antagonize muscarinic receptors. They inhibit reflex cholinergic bronchoconstriction and do not significantly block the direct effects of inflammatory mediators such as histamine and leukotrienes on bronchial smooth muscle and vessels. When given by inhalation, anticholinergics produce bronchodilation by competitively inhibiting cholinergic receptors in bronchial

smooth muscle. This activity blocks acetylcholine with the net effect being a reduction in cyclic guanosine monophosphate (cGMP) that normally acts to constrict bronchial smooth muscle. Anticholinergic drugs usually are less effective as bronchodilators in asthmatic subjects than β adrenergic agonists. Nevertheless, they may have an additive effect with β adrenergic agonists.

Atropine reduces mucociliary clearance in normal subjects and in patients with asthma and chronic bronchitis, but the quaternary derivative, ipratropium bromide, even when given in high doses, has no such detectable effect either on normal subjects or in patients with airway disease (Pavia et al. 1980).

Ipratropium bromide has been shown to decrease the effectiveness of voluntary cough on clearing mucus from the airways, which may affect its role in the treatment of patients who have excessive mucus production. Ipratropium has a slower onset of action and a more prolonged bronchodilator effect compared with standard β_2 -agonists and has been considered to be less suitable for use on an as needed basis for immediate relief of bronchospasm.

The lack of systemic absorption of ipratropium greatly diminishes the anticholinergic side effects such as blurred vision, urinary retention, nausea, and tachycardia associated with atropine. A significant unwanted effect of inhaled ipratropium bromide is dryness of mouth and throat, bitter taste, cough and nausea. Nebulized ipratropium bromide may precipitate glaucoma in elderly patients because of its direct mydriatic effect on the eye. During sleep, ipratropium also has been shown to improve arterial oxygen saturation and sleep quality.

Tiotropium bromide is a long acting quaternary anticholinergic agent. Tiotropium in human lungs shows approximately 10 fold more potency than ipratropium and protects against cholinergic bronchoconstriction for greater than 24 h.

Methylxanthines

Methylxanthines such as theophylline are related to caffeine and have been used to treat asthma since 1930. The methylxanthines may produce bronchodilation through numerous mechanisms, including,

- inhibition of phosphodiesterase, thereby increasing cAMP levels
- inhibition of calcium ion influx into smooth muscle
- prostaglandin antagonism
- stimulation of endogenous catecholamines
- adenosine receptor antagonism
- Inhibition of release of mediators from mast cells and leukocytes.

Theophylline inhibits release of mediators from mast cells, increases mucociliary clearance, and prevents the development of micro vascular leakiness, as would an

“anti-inflammatory” drug (Persson and Draco 1988). Theophylline also inhibits some functions of T lymphocytes, which may be relevant to control of chronic inflammation of the airway.

For nocturnal asthma, a single dose of slow release theophylline at bedtime often is effective. This has been demonstrated to reduce overnight declines in FEV₁ and morning respiratory symptoms. Taken alone it increases exercise tolerance without improving spirometry tests.

Other theophylline salts, such as choline theophyllinate, offer no advantages over theophylline. The ethylenediamine component of aminophylline has been implicated in allergic reactions. Some derivatives such as acepiophylline, diprophyl-line, and proxophylline, are less effective than theophylline (Weinberger 1984). The most common adverse effects are headache, nausea and vomiting, abdominal discomfort, and restlessness.

Anti-inflammatory drugs

Although the type of inflammatory responses may differ among diseases, inflammation is a common denominator of several lung diseases. Anti-inflammatory drugs suppress the inflammatory response by inhibiting infiltration and activation of inflammatory cells as well as their synthesis or release of mediators or effects of inflammatory mediators themselves.

Corticosteroids

Since asthma is viewed as a chronic inflammatory disease and inhaled corticosteroids are known to have low toxicity, they may be considered as first line therapy (Barns 1989). Prednisolone and dexamethasone were effective when they were given systematically to treat asthma but they had no anti-asthmatic activity when they were given by inhalation. Other corticosteroids e.g. beclomethasone dipropionate (BDP), betamethasone and budesonide, were effective in treating asthma when given by inhalation. The antiasthmatic potency of an inhaled steroid is approximately proportional to its potency as an anti-inflammatory agent.

Corticosteroids inhibit the release of arachidonic acid metabolites and platelet activating factor (PAF) from lungs and macrophages by enhancing the production of proteins called lipocortin. Thereby they inhibit the formation of prostaglandins and leukotrienes. These effects occur because of ability of steroid—receptor complex to be transported to the nucleus, where it initiates DNA transcription of specific mRNAs. Corticosteroids potentially inhibit the accumulation of neutrophils, inhibit secretion of human pulmonary macrophages of leukotrienes and prostaglan-

dins, inhibit formation of interleukins (ILs) such as IL-1, IL-2, IL-3 and IL-5, inhibit degranulation and adherence of eosinophils, reduce number of circulating T lymphocytes and formation of an IgE binding suppressive factor. Steroids prevent and reverse the increase in vascular permeability due to inflammatory mediators and may therefore lead to resolution of airway edema. Corticosteroids remain the most effective therapy available for asthma but the legitimate fear of their adverse effects makes using them difficult. Steroids potentiate the effects of β adrenergic agonists on bronchial smooth muscle (Barns 1989). Methylprednisolone is given intravenously to patients with severe acute asthma. Inhaled steroids have no proven value in the management of acute asthma. Patients with chronic bronchitis occasionally respond to steroids, possibly because some have an element of undiagnosed asthma.

Corticosteroids inhibit release of ACTH and secretion of cortisol by a negative feed back effect on the pituitary gland. Adverse effects of corticosteroids include fluid retention, increased cell mass, increased appetite, weight gain, osteoporosis, capillary fragility, hypertension, peptic ulceration, diabetes, cataract, and psychosis (Dajani et al. 1981).

Anti-leukotrienes

Leukotrienes possess potent pro-inflammatory actions resulting in increased vascular permeability, mucus secretion and bronchial hyperresponsiveness. They are derived from the 5-lipoxygenase pathways in mast cells, eosinophils and macrophages. Anti-leukotrienes improve lung function and diminish symptoms, exacerbation rate and the need for rescue bronchodilator. These are drugs of choice in case of aspirin induced asthma, in which patients have high LTE_4 levels in urine and nasal secretions and even higher after taking aspirin (Christie et al. 1992).

Leukotriene modifiers are drugs that modify the response of these mediators of inflammation by one of the four ways (Drazen 1997).

a) Cysteinyl LT receptor inhibitors

C-LTs promote eosinophil influx, bronchospasm and mucus hypersecretion, all are considered hallmarks of asthma. C-LT receptor inhibitors antagonize or inhibit leukotrienes predominantly LTD_4 . These agents inhibit phospholipases, prostaglandins, leukotrienes, and IL-1 synthesis. Pranlukast and Irelukast belong to this class (Drazen 1997; Floreani and Rennard 1999).

b) 5-lipoxygenase inhibitors

They prevent the formation of leukotrienes by blocking a 5-lipoxygenase pathway in their synthesis. Zileuton, ZD-2138, ABt-761 belongs to this class (Floreani and Rennard 1999).

c) 5-lipoxygenase activating protein (FLAP) inhibitors

MK-0591 and MK-886 attenuated the early and late asthmatic response following antigen challenge but not the attendant increase in airway responsiveness to spasmogens (Diamant et al. 1995).

d) Leukotrienes receptor antagonists

Montelukast, Zafirlukast, Pranlukast are selective and high affinity LT_1 antagonists (Adcock and Matthews 1998).

Zileuton has shown efficacy in exercise-induced asthma, aspirin induced bronchospasm and following chronic administration, an improvement in pulmonary function (FEV1) and a reduction in oral and inhaled corticosteroid use (Tamaoki et al. 1997). Furthermore, in a small study, zileuton attenuated both airway and blood eosinophilia in nocturnal asthmatics (Wenzel et al. 1995).

Zafirlukast has been demonstrated to attenuate the acute airway obstructive response to allergen and exercise challenge and to improve chronic asthma control both objectively (FEV1, nocturnal awakenings, β -agonist use) and subjectively.

Montelukast has been shown to block the early and late response to allergen challenge following single dosing, to improve FEV1 in both children (6–14 years) and adults and to protect against the development of exercise induced bronchoconstriction in both children and adults. Tolerance to the bronchoprotective effects of montelukast in attenuating exercise-induced bronchospasm does not develop following at least 12 weeks of therapy.

Pranlukast increases FEV1 within 1 h of dosing, improves patient summary symptom and nighttime asthma scores and reduces the use of rescue bronchodilators. In patients with moderate persistent asthma, it prevents exacerbations of asthma during reduction of high dose inhaled corticosteroids therapy (Tamaoki et al. 1997).

Mediator release inhibitors

Cromolyn sodium

Cromolyn Sodium (Sodium cromoglycate) is a derivative of khellin, an Egyptian herbal remedy. Cromolyn inhibited the release of mediators by allergen in passively sensitized animal and human lung preparations (Cox 1967). Cromolyn was classified as mast cell stabilizer. Cromolyn has variable inhibitory actions on other inflammatory cells including macrophages and eosinophils that may participate in allergic inflammation. In vivo cromolyn can block both the early response that may be mediated by mast cells to allergens and the late response and bronchial hyperresponsiveness (Cockcroft and Murdock 1987). Cromolyn Sodium is used for prophylactic treatment and consequently needs to be taken regularly. It is the first choice anti-

inflammatory drug for children because it has few adverse effects (Bernstein 1985). Cromolyn sodium is classified as an antiallergic drug because it appears to have a specific effect on allergy based inflammation. Several other drugs also may be included in this category.

Nedocromil sodium is a new drug used for prophylaxis. It has a similar pharmacologic profile of activity to cromolyn, is more potent in various tests, and may have a longer duration of action. Ketotifen also is described as a drug to be used for prophylaxis against asthma.

Newer targets in asthma therapy

The current pharmacotherapeutic approaches to asthma have several limitations. First, there is no known asthma cure and little evidence that prevention is possible in susceptible persons. Hence, patients continue to be at risk of symptoms

and exacerbations. Mortality remains a severe problem. Finally, the medications have adverse effects. There is even some evidence, albeit conflicting, that cataract formation, osteoporosis and growth impairment, as associated with systemic glucocorticoids, may arise from topical steroids, depending on dosages used. New inhalation devices and new generation beta-agonists are available. At the same time, new understanding of the molecular pathology of asthma has identified several novel therapeutic targets. Agents being tested in early phase clinical trials include antagonists of IgE, cytokines, adhesion molecules and transcription factors.

TXA₂ inhibitors

TXA₂ is a potent bronchoconstrictor, mucus producer and blood vessel permeability inducer and causes airway hyper responsiveness. Serabonast, domitroban and ozagrel are the

Table 1 Bronchodilators

Sr. No	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Adhatoda vasica</i> Nees	Leaves, Roots	Alkaloids	Paliwa et al. 2000
2.	<i>Albizzia lebbeck</i> (<i>Sareesha_rakat</i>)	Stem bark/Aqueous	Saponins	Tripathi and Das 1977
3.	<i>Alstonia scholaris</i>	Leaves/Ethanol	Ditamine, Echitamine and Echitenines	Channa et al. 2003
4.	<i>Artemisia caerulescens</i>	Aerial parts/Butanol	Quercetin, isorhamnetin	Moran et al. 1989
5.	<i>Belamcanda chinensis</i>	Leaves/Ethanol	Tectorigenin	Singh and Agrawal 1990
6.	<i>Benincasa hispida</i>	Fruits/Methanol	Triterpenes, Glycosides, Sterols	Kumar and Ramu 2002
7.	<i>Cissampelos sympodialis</i>	Leaves and root bark/Aqueous	Warifteine, α -bisbenzylisoquinoline alkaloid	Thomas et al. 1995, 1997; Cortes et al. 1995
8.	<i>Clerodendron serratum</i>	Stem bark/Aqueous	Phenolic glycoside	Gupta 1968; Gupta and Tripathi 1973
9.	<i>Coleus forskohlii</i>	Roots	Forskolin (Diterpenoid)	Marone et al. 1987
10.	<i>Elaeocarpus sphaericus</i>	Fruits/Aqueous, Pet-ether, Benzene, Acetone and ethanol	Glycoside, Steroids, Alkaloid, Flavanoids	Singh et al. 2000
11.	<i>Galphimia glauca</i>	Aerial/Alcohol extract/Ethyl-acetate	Tetragalloylquinic acid, Quercetin	Campos et al. 2001
12.	<i>Gardenia latifolia</i>	Bark	Saponins	Gupta 1974
13.	<i>Ginko biloba</i>	Leaves	Ginkgolides	Puglisi et al. 1988
14.	<i>Mikania glomerata</i>	Leaves/Aqueous, hydroalcohol	Coumarin	Soares de Moura et al. 2002
15.	<i>Lepidium sativum</i>	Seeds/Ethanol fractions	Alkaloids, Flavonoids	Mali et al. 2008
16.	<i>Ocimum sanctum</i>	Leaves/Ethanol	Myrcenol, Nerol, Eugenol	Singh and Agrawal 1991
17.	<i>Passiflora incarnata</i>	Leaves/Methanol	Alkaloids	Dhawan et al. 2003
18.	<i>Pavetta crassipes</i>	Leaves/Aqueous	Flavanoids, tannins, anthraquinones	Amos et al. 1998
19.	<i>Picrorrhiza kurroa</i>	Roots/	Androsin	Stuppner et al. 1991
20.	<i>Sarcostemma brevistigma</i>	Twigs/Alkaloidal fraction	Bregenin	Saraf and Patwardhan 1988b
21.	<i>Tephrosia purpurea</i>	Aerial parts/Ethanol extract	Flavanoids, Tephrosin	Gokhale et al. 2000
22.	<i>Tylophora indica</i>	Leaves/Alkaloidal fraction	Tylophorine	Nayampalli et al. 1986
23.	<i>Vitex negundo</i>	Leaves/Ethanol	Casticin, isoorientin Chrysophenol D, Luteolin	Nair and Saraf 1995
24.	<i>Rosmarinus officinalis</i>	Shrub/Aqueous	Caffeic acid (CA) and Rosmarinic acid	Aqel 1991
25.	<i>Ephedra sinica</i>	Stems	Ephedrine	Akiba et al. 1979
26.	<i>Gleditsia sinensis</i> Lam.	Leaves/Decoctions		Dai et al. 2002

examples of these TXA₂ synthetase inhibitors. Ozagrel reduced cough sensitivity to capsaicin and bronchoconstriction due to acetaldehyde. TXA₂ antagonists *BAÿu3405* produced a modest decrease in airways responsiveness to methacholine following 2 weeks treatment in asthmatics.

Tachykinin receptor antagonists

The first nonpeptide tachykinin receptor antagonist was *CP-96345*, which is a potent NK₁ receptor antagonist. *SR*

48968, *GR 159897* and *SR 144190* are selective nonpeptide NK₂ receptor antagonists. *SR 142801* and *SB 223412* are selective NK-3 receptor antagonists.

Tryptase inhibitors

Tryptase inhibitors inhibit both early and late reactions. *APC-366* inhibited antigen induced late phase response and bronchial hyperresponsiveness to carbachol in sheep. Lactoferrin disrupts the quaternary structure

Table 2 Mast cell stabilizers

Sr.No.	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Achyranthes aspera</i>	Aerial parts/Aqueous	Oleanolic acid	Agrawal and Mehta 2005
2.	<i>Albizia lebbbeck</i>	Stem bark/Aqueous	Saponins	Tripathi et al. 1979
3.	<i>Allium cepa</i>	Bulbs/Juice	α and β unsaturated Thiosulphinates	Johri et al. 1985
4.	<i>Aquillaria agallocha</i>	Stem/Aqueous extract	Triterpenoids	Kim et al. 1997
5.	<i>Azadirachta indica</i>	Leaves/Juice	Nimbin, nimbinine, Nimbandiol, quercetin	Acharya et al. 2003
6.	<i>Bacopa monniera</i>	Leaves/Ethanol	Bacosides, Alkaloids, Glycosides	Samiulla et al. 2001
7.	<i>Bidens parviflora</i>	Aerial parts	Glycosides	Wang et al. 2001
8.	<i>Calotropis procera</i>	Latex	α -amyrin, β -amyrin calotropin (Triterpenoid)	Kumar and Basu 1994
9.	<i>Cassia alata</i>	Leaves/Ethanol	Anthraquinones, Flavanoids	Palanichamy et al. 1991
10.	<i>Cassia obtusifolia</i>	Seeds/Glycosidal fraction	Anthraquinones, Betulinic acid	Kitanaka et al. 1998
11.	<i>Cassia torosa</i>	Seeds	Gentiobiosides	Kanno et al. 1999
12.	<i>Cedrus deodara</i>	Wood oil	Himacholol	Shinde et al. 1999
13.	<i>Citrus unshiu</i>	Peels/	Flavanoids	Kim et al. 1999
14.	<i>Clerodendron serratum</i>	Bark/Aqueous	Phenolic glycoside	Gupta 1968
15.	<i>Cnidium monnieri</i>	Fruits/Ethanol	Osthof	Chen et al. 1988
16.	<i>Coleus forskohlii</i>	Roots	Forskolin (diterpenoid)	Marone et al. 1987
17.	<i>Crinum glaucum</i>	Leaves/Aqueous	Alkaloids, lycorine, crinamine	Okpo and Adeyemi 2002
18.	<i>Curcuma longa</i>	Rhizome	Tumerones, curcuminoids	Ammon and Wahl 1991
19.	<i>Drymaria cordata</i>	Methanol extracts		Mukherjee et al. 1997
20.	<i>Elaeocarpus sphaericus</i>	Fruits/Aqueous, Pet-ether, Benzene, Acetone and ethanol	Glycoside, Steroids, Alkaloid, Flavanoids	Singh et al. 2000
21.	<i>Gleditsia sinensis</i>	Fruits/Ethanol	Saponins	Dai et al. 2002
22.	<i>Impatiens textori</i>	Flowers/Ethanol	Apigenin, uteolin, chrysoeriol	Ishiguro et al. 2000
23.	<i>Inula racemosa</i>	Roots/Alcohol	Inulolide-a new Sesquiterpene lactone	Srivastava et al. 1999
24.	<i>Magnolia officinalis</i>	Bark/Aqueous	Honokiol, Magnolol	Shin et al. 2001b
25.	<i>Mentha piperita</i>	Leaves	Flavanoidal glycosides	Inoue et al. 2002
26.	<i>Ocimum sanctum</i>	Leaves/Aqueous	Myrcenol, Nerol, Eugenol	Sen 1993
27.	<i>Picrorrhiza kurroa</i>	Roots/	Androsin	Stuppner et al. 1991
28.	<i>Solanum xanthocarpum</i>	Roots/Alkaloidal fraction	Solasodine	Chitravanshi et al. 1990
29.	<i>Striga orobanchioids</i>	Aerial parts/Ethanol		Harish et al. 2001
30.	<i>Tephrosia purpurea</i>	Aerial parts/Ethanol extract	Flavanoids, Tephrosin	Gokhale et al. 2000
31.	<i>Terminalia chebula</i>	Fruits/Aqueous	Ellagic acid, Tannins Chebulagic acid	Shin et al. 2001a
32.	<i>Tinospora cordifolia</i>	Stem/Aqueous	Tinosporin	Nayampalli et al. 1986
33.	<i>Tylophora asthmatica</i>	Leaves/Alkaloidal	Tylophorine	Geetha et al. 1981
34.	<i>Vitex negundo</i>	Leaves/Ethanol	Casticin, isoorientin Chrysophenol D, Luteolin	Nair et al. 1994

of tryptase, also attenuates antigen induced late response and bronchial hyperresponsiveness in allergic sheep.

Cytokine inhibitors

One of the novel approaches for the treatment of asthma is to target cytokines and develop cytokine modulators as drugs. Two humanized anti-IL-5 monoclonal antibodies, *Sch-55700* and *SB-240563* reduced blood eosinophil count for several weeks and prevented eosinophils recruitment into the airways after allergen challenge in asthmatic patients. IL-5 signaling inhibitor *GCC-AP0341* inhibited IL-5 mediated survival of eosinophils. IL-4 receptor antibodies inhibited allergen induced airway hyperresponsiveness, goblet cell metaplasia and pulmonary eosinophilia in a murine model.

Chemokine inhibitors

A variety of chemokines, one of which is the chemo-attractant eotaxin, are secreted by inflamed lung tissue thereby attracting eosinophils. Eotaxin receptor blockers are being investigated, as eosinophils are believed to be major contributors to the pulmonary damage seen in asthma. Monoclonal antibody (7B11) for human CCR₃ has shown to completely block the binding and signaling of the known CCR₃ ligands, thus blocking the chemotactic response of human eosinophils to all chemokines.

Adhesion molecule antagonists

Interactions of eosinophils with intra cellular adhesion molecule-1 (ICAM-1) are thought to be necessary for eosinophils recruitment into airways. Antibodies to ICAM-1

Table 3 Anti-allergics

Sr. No.	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Adhatoda vasica</i>	Leaves/Methanol	Vasicinol, vasicine	Muller et al. 1993; Kumar Suresh 1979
2.	<i>Albizia lebbbeck</i>	Stem bark/Aqueous	Saponins	Baruah et al. 1997; Suresh et al. 1981
3.	<i>Alisma orientale</i>	Rhizomes/Aqueous, Methanol	Alisol B monoacetate, Alismaketones-B 23-acetate and -C 23-acetate	Kubo et al. 1997
4.	<i>Aquillaria agallocha</i>	Stem/Aqueous extract	Triterpenoids	Kim et al. 1997
5.	<i>Asiasarum sieboldi</i>	Roots/Methanol	Methyleugenol, gamma-asarone, Elemicin, Asarinin	Hashimoto et al. 1994
6.	<i>Camellia sinensis</i>	Leaves	Flavanoids	Suzuki et al. 2000
7.	<i>Centipeda minima</i>	Aerial parts	Flavanoids, Pseudoquinolide, sesquiterpene lactones	Wu et al. 1985
8.	<i>Citrus unshiu</i>	Peels/	Flavanoids	Kim et al. 1999
9.	<i>Cnidium monnieri</i>	Fruits/Ethanol	Osthol	Matsuda et al. 2002
10.	<i>Crinum glaucum</i>	Leaves/Aqueous	Alkaloids, lycorine, crinamine	Okpo and Adeyemi 2002
11.	<i>Curcuma longa</i>	Rhizomes	Curcumin and tetrahydrocurcumin	Suzuki et al. 2000
12.	<i>Dalbergia odorifera</i>	Heart Wood	Flavanoids, Tannins	Chan et al. 1998
13.	<i>Desmodium adscendens</i>	Aqueous	Triterpenoid Saponin	Addy 1989
14.	<i>Galphimia glauca</i>	Aerial/Alcohol extract/ Ethyl-acetate	Tetragalloylquinic acid, Quercetin	Neszmelyi et al. 1993
15.	<i>Ginkgo biloba</i>	Leaves	Ginkgolides	Touvay et al. 1985
16.	<i>Gleditsia sinensis</i>	Fruits/Ethanol	Saponins	Dai et al. 2002
17.	<i>Hydrangea macrophylla</i>	Leaves	Glycosides	Matsuda et al. 1999
18.	<i>Inula racemosa</i>	Roots/Alcohol	Inulolide-a new Sesquiterpene lactone	Srivastava et al. 1999
19.	<i>Magnolia officinalis</i>	Bark/Aqueous	Honokiol, Magnolol	Shin et al. 2001b
20.	<i>Sarcostemma brevistigma</i>	Twigs/Alkaloidal fraction	Bregenin	Saraf and Patwardhan 1988a
21.	<i>Solanum xanthocarpum</i>	Roots/Alkaloidal fraction	Solasodine	Chitravanshi et al. 1990
22.	<i>Terminalia chebula</i>	Fruits/Aqueous	Ellagic acid, Tannins Chebulagic acid	Shin et al. 2001a
23.	<i>Vitex negundo</i>	Leaves/Ethanol	Casticin, isoorientin Chrysophenol D, Luteolin	Nair and Saraf 1995

Table 4 Anti-inflammatory agents

Sr.No.	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Asystasia gangetica</i>	Leaves/Methanol, Ethyl Acetate	Isoflavone glycoside, dalhorinin	Akah et al. 2003
2.	<i>Aloe vera</i> Tourn.ex Linn. (Liliaceae)	Leaves/Aqueous, Chloroform and ethanol	Anthraquinones, sterols, saponins and carbohydrate	Vázquez et al. 1996
3.	<i>Bryonia laciniosa</i>	Leaves/chloroform extract	Flavonoids	Gupta et al. 2003
4.	<i>Calotropis procera</i>	Latex	α -amyrin, β -amyrin calotropin (Triterpenoid)	Kumar and Basu 1994
5.	<i>Cinnamomun Zeylanicum</i>	oil	Eugenol, cinnamic aldehyde and α -terpeniol.	
6.	<i>Curcuma longa</i>	Rhizomes	Tumerones, curcuminoids	Ammon and Wahl 1991
7.	<i>Dalbergia odorifera</i>	Heart Wood	Flavanoids, Tannins	Chan et al. 1998
8.	<i>Elaeocarpus sphaericus</i>	Fruits/Aqueous, Pet-ether, Benzene, Acetone and ethanol	Glycoside, Steroids, Alkaloid, Flavanoids	Singh et al. 2000
10.	<i>Nelsonia canescens</i>	Leaf/ethanol extract	Flavonoids	Owoyele et al. 2005
11.	<i>Indigofera tinctoria</i>	Whole plant/methanol	Polyphenols	Oli et al. 2005
12.	<i>Butea frondosa</i> Koen.	Leaves/Aqueous	Flavonoid, glycosides, proteins and amino acids.	Mengi and Deshpande 1999
13.	<i>Ocimum sanctum</i>	Leaves/Aqueous	Myrcenol, Nerol, Eugenol	Singh and Agrawal 1991
14.	<i>Ophiopogon japonicus</i>	Root/aquoes extract	Ruscogenin and ophiopogonin D	Kou et al. 2005
15.	<i>Pavetta crassipes</i>	Leaves/Aqueous	Flavanoids, tannins, anthraquinones	Amos et al. 1998
16.	<i>Tylophora asthmatica</i>	Leaves/Alkaloidal	Tylophorine	Manez et al. 1990

Table 5 Anti-spasmodic agents

Sr. No.	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Aegle marmelos</i>	Leaves/Ethanol	Aegelin, Aegelemine, Aegeline	Arul et al. 2004
2.	<i>Asiasarum sieboldi</i>	Roots/Methanol	Methyleugenol, gamma-asarone, Elemicin, Asarinin	Hashimoto et al. 1994
3.	<i>Asystasia gangetica</i>	Leaves/Methanol, Ethyl acetate	Isoflavone glycoside, dalhorinin	Akah et al. 2003
4.	<i>Bacopa monniera</i>	Leaves/Ethanol	Bacosides, Alkaloids, Glycosides	Dar and Channa 1997; Channa et al. 2003
5.	<i>Belamcanda chinensis</i>	Leaves/Ethanol	Tectorigenin	Singh and Agrawal 1990
6.	<i>Cissampelos glaberrima</i>	Leaves, Root Bark/Aqueous	Warifteine, α -bisbenzylisoquinoline alkaloid	Thomas et al. 1995; Cortes et al. 1995
7.	<i>Clerodendron serratum</i>	Stem bark/Aqueous	Phenolic glycoside	Gupta 1968
8.	<i>Cnidium monnieri</i>	Fruits/Ethanol	Osthol	Chen et al. 1988
9.	<i>Coleus forskohlii</i>	Roots	Forskolin (diterpenoid)	Marone et al. 1987
10.	<i>Crinum glaucum</i>	Leaves/Aqueous	Alkaloids, lycorine, crinamine	Okpo and Adeyemi 2002
11.	<i>Drymis winteri</i>	Bark	Terpene	Sayah et al. 1998
12.	<i>Ferula ovina</i>	Aerial parts/Ethanol	Carvacrol, alpha-pinene, geranyl isovalerate and geranyl propionate	Khalil et al. 1990
13.	<i>Ferula sinica</i>	Roots/Ethanol	Resins	Aqel et al. 1991
14.	<i>Pavetta crassipes</i>	Leaves/Aqueous	Flavanoids, tannins, anthraquinones	Amos et al. 1998
15.	<i>Saussurea leppa</i>	Alkaloidal fraction	Sesquiterpene lactone, Terpenoids	Dutta et al. 1968
16.	<i>Striga orobanchioids</i>	Aerial parts/Ethanol	??	Harish et al. 2001
17.	<i>Thymus vulgaris</i>	Ethanol	Flavanones	Meister et al. 1999
18.	<i>Tylophora asthmatica</i>	Leaves/Alkaloidal	Tylophorine	Haranath and Shyamalakumari 1975; Udapa et al. 1991

blocked both eosinophils recruitment into the airways in the monkey model of asthma and importantly the increase in airway reactivity associated with allergen challenge

Phosphodiesterase inhibitors

Considerable interest has been generated in the potential utility of isoenzyme-selective inhibitors of cyclic nucleotide Phosphodiesterase (PDE) in the treatment of asthma and other inflammatory disorders. The scientific foundation for this interest is based upon two fundamental principles. First, inhibition of PDE activity increases the cellular content of two key second messengers, cAMP and cGMP, thereby activating specific protein phosphorylation cascades that elicit a variety of functional responses. Increases in cAMP content suppress a broad array of functions in inflammatory and immune cells. Both cAMP and cGMP mediate bronchodilation. PDE3 inhibitor enoxamine was shown to decrease lung resistance and increase compliance in patients with decompensated chronic pulmonary disease. Benzafranine administered to normal volunteers by inhalation produced bronchodilation. Zaprinas is PDE5 inhibitor; it reduced exercise-induced bronchoconstriction but not histamine-induced bronchoconstriction. Most of the

work now is focused on selectively targeting PDE₄, primarily because inhibitors of this isoenzyme family have a notably appealing therapeutic profile; broad-spectrum anti-inflammatory activity coupled with additional bronchodilatory and neuromodulatory action. Rolipram, *LAS-31025*, *RP-73401* and denbufylline are selective PDE₄ inhibitors. *SB 207499*, *VII294A*, *CP-220* and roflumilast are PDE₄ inhibitors with less gastrointestinal side effects.

Endothelin modulators

There are two approaches for ET-1 directed therapeutics- (1) Inhibitors of endothelin-converting enzyme (ECE), which mediates the synthesis of ET-1 from its precursor; (2) Receptor antagonists of the effects of ET-1 at the end organ level. These agents reverse and/or prevent the increase in pulmonary artery pressure and vascular remodeling elicited by acute or chronic hypoxia. Examples are BQ-123, SB-217242 and bosentan.

Herbal drugs in bronchial asthma

Many Ayurvedic plants have been described to be useful in the treatment of various bronchial disorders including

Table 6 Miscellaneous agents

Sr No	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
Lipoxygenase inhibitors				
1.	<i>Allium cepa</i>	Bulbs/Juice	α and β unsaturated Thiosulphinates	Bayer et al. 1989
2.	<i>Boswellia serrata</i>	Gum resin/Ethanol	Boswellic acid	Ammon et al. 1991
3.	<i>Coleus forskohlii</i>	Roots	Forskolin (diterpenoid)	Marone et al. 1987
4.	<i>Proustia pyrifolia</i>	Whole plant/methanol	B-sitosterol, quercetin and dihydroquercetin	Delparte et al. 2005
Platelet Activating Factor (PAF) inhibitors				
1.	<i>Allium cepa</i>	Bulbs/Juice	α and β unsaturated Thiosulphinates	Dorsch et al. 1987
2.	<i>Galphimia glauca</i>	Aerial/Alcohol extract/ Ethyl-acetate	Tetragalloylquinic acid, Quercetin	Neszmelyi et al. 1993
3.	<i>Impatiens textori</i>	Flowers/Ethanol	Apigenin, uteolin, chrysoeriol	Ueda et al. 2003
4.	<i>Picrorrhiza kurroa</i>	Roots/	Androsin	Stuppner et al. 1991
Cyclooxygenase inhibitor				
1.	<i>Allium cepa</i>	Bulbs/Juice	α and β unsaturated Thiosulphinates	Bayer et al. 1989
2.	<i>Magnolia obovate</i>	Stem bark	Magnolol and honokiol	Lee et al. 2005
3.	<i>Crataegus pinnatifida</i>	Fruit	Flavanoids	Kao et al. 2005
Interleukins (ILs) biosynthesis inhibitors				
1.	<i>Calocedrus formosana</i>	Bark/alcohol	Sugiol	Chao et al. 2005
2.	<i>Nidularium procerum</i>	Leaves/Aqueous extract		Vieira-de-Abreu et al. 2005
3.	<i>Waltheria indica</i>	Whole plant/	flavanoids	Rao et al. 2005
4.	<i>Mahonia aquifolium</i> Nutt. Berberidaceae	Stem bark/hydroalcohol extract	Polysaccharides Protoberberine and bisbenzylisoquinoline (BBI) alkaloids berbamine, tetrandrine	Kostalova et al. 2001
Leukotriene biosynthesis inhibitors				
1.	<i>Nigella sativa</i>	Seeds oil	Thymoquinone (TQ)	El Gazzar et al. 2006

Table 7 Antianaphylactic drugs

Sr. No.	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Xanthii fructus</i>	Whole plant/Aqueous	Saponin, flavones, Caffeic acid, 1,4-dicaffeoylquinic acid, sesquiterpene lactones	Hong et al. 2003
2.	<i>Lycopus lucidus</i>	Whole plant/Aqueous	Betulinic acid, pentacyclic Triterpenes	Shin et al. 2005
3.	<i>Poncirus trifoliata</i>	Fruit/Aqueous	Flavonoids	Kim et al. 1999
4.	<i>Trichopus zeylanicus</i>	Leaves/butanol	Lipoprotein/Glycolipoprotein	Subramoniam et al. 1999
5.	<i>Cryptotympana atrata</i>	Whole plant/Aqueous	oleanolic acid	Shin et al. 1999
6.	<i>Striga orobanchioides</i>	Whole plant/Aqueous, ethanolic	Flavonoids, apigenin and luteolin	Harish et al. 2001
7.	<i>Crinum glaucum</i>	Bulbs/Aqueous	Alkaloids	Okpo and Adeyemi 2002
8.	<i>Acanthopanax senticosus</i>	Stem/Aqueous	Acanthoside A, B & C, Chiisanoside, Senticoside, Saponin, flavones,	Yi et al. 2002
9.	<i>Syzygium aromaticum</i>	Flower bud/Aqueous	Phenols	Kim et al. 1998
10.	<i>Terminalia chebula</i>	Fruit/Aqueous	Tannins	Shin et al. 2001a
11.	<i>Vitex rotundifolia</i>	Fruit/Aqueous	Flavonoids	Shin et al. 2000

bronchial asthma (Kumar Suresh 1979). The use of medicinal plants and natural products increased dramatically in the last two decades in all over the world. More than 400 medicinal plant species have been used ethanopharmacologically and traditionally to treat the symptoms of asthmatic and allergic disorders worldwide.

Classification of anti asthmatic herbs based on mechanism of action

Some herbal alternatives employed in asthma are proven to provide symptomatic relief and assist in the inhibition of disease development as well. These herbs therefore have multifaceted roles to play in the management of asthma

suggesting different sites of action within the body. Based on the possible mechanism of action reported, plant anti-asthmatics may be classified as shown in tables (Tables 1, 2, 3, 4, 5, 6, 7 and 8).

Conclusion

Herbal approaches have regained their popularity, with their efficacy and safety aspects being supported by controlled clinical studies. The herbal approaches have offered effective mast cell stabilizers like sodium cromolyn and sodium cromoglycate developed from khellin and anti-leukotriene products like—boswellic acids. Ongoing re-

Table 8 Immunomodulatory drugs

Sr. No.	Name of plant	Part used/extract/fraction	Major chemical constituent(s)	Reference
1.	<i>Picrorhiza scrophulariiflora</i>	Rhizomes/Pet. ether, Diethyl ether and methanol	Apocynin, androsin and picroside II	Smit et al. 2000
2.	<i>Trichilia glabra</i>	Leaf/Aqueous	Polysaccharides	Benancia et al. 2000
3.	<i>Cedrela tubiflora</i>	Leaf/Aqueous	Gallic acid, polysaccharides	Benancia et al. 1995
4.	<i>Ipomoea carnea</i>	Leaf/Aqueous	Nortropane alkaloids, calystegines β_2	Hueza et al. 2003
5.	<i>Withania somnifera</i>	Coded extracts		Rasool and Varalakshmi 2006
6.	<i>Clausena excauata</i>	Wood/Aqueous	Phenolic compounds, furanocoumarins, flavanoids and carbazole alkaloid	Manosroi et al. 2005
7.	<i>Magnifera indica</i>	Bark/Alcohol, ether	Magniferin	Makare et al. 2001
8.	<i>Cleome viscosa</i>	Aerial parts/Aqueous, ethanolic	Alkaloids, saponins	Tiwari et al. 2003
9.	<i>Plantago ovata</i>	Seeds/Aqueous	Polysaccharides glycosides	Rezaeipoor et al. 2000
10.	<i>Typhae angustifolia</i>	Pollen/ethanol	Phenolic compounds, flavones Triterpenes And β -sitosterol	Qin and Sun 2005
11.	<i>Angelica sinensis</i>	Roots/Aqueous and ethanolic	Polysaccharides	Yang et al. 2006
12.	<i>Boerhaavia diffusa</i>	Roots/ethanol	Alkaloids	Mungantiwar et al. 1999
13.	<i>Tephrosia purpurea</i>	Aerial parts/Ethanol	Flavonoids	Damre et al. 2003

search worldwide has provided valuable clues regarding the precise mechanism of action of these herbal alternatives and these herbs, have shown interesting results in various target specific biological activities such as bronchodilation, mast cell stabilization, anti-anaphylactic, anti-inflammatory, anti-spasmodic, anti-allergic, immunomodulatory and inhibition of mediators viz., leukotrienes, lipoxygenase, cyclooxygenase, platelet activating, phosphodiesterase and cytokine, in the treatment of asthma.

Some herbal alternatives employed in these traditions are proven to provide symptomatic relief and assist in the inhibition of disease development as well. In nutshell, attempt should be made to develop polyherbal formulations which contain various herbs acting at particular sites of the pathophysiological cascade of asthma for prophylaxis as well as for the treatment of asthma and subsequent clinical studies on them.

References

- Acharya SB, Yanpallewar SU, Singh RK (2003) A preliminary study on the effect of *Azadirachta indica* on bronchial smooth muscles and mast cells. *J Nat Rem* 3:78–82
- Adcock IM, Matthews JC (1998) New drugs for asthma. *Drug Discov Today* 3:395–419
- Addy ME (1989) Several chromatographically distinctive fractions of *Desmodium adscendens* inhibit smooth muscle contractions. *Int J Crude Drug Res* 27:81–91
- Agrawal BB, Mehta AA (2005) Phyto-pharmacological investigation of *Moringa oleifera* and *Achyranthus aspera* for their anti-asthmatic activity. Ph.D. thesis, Gujarat University
- Akah PA, Ezike AC, Nwafor SV, Okoli CO, Enwerem NM (2003) Evaluation of the anti-asthmatic property of *Asystasia gangetica* leaf extracts. *J Ethnopharmacol* 89:25–36
- Akiba K, Onodera K, Kisara K, Fujikura H (1979) Interaction of d-pseudoephedrine with water soluble extracts of *Platycodi radix* on acute toxicity. *Nippon Yakurigaku Zasshi* 75:201–206
- Ammon HP, Mack T, Singh GB, Safayhi H (1991) Inhibition of leukotriene B₄ formation in rat peritoneal neutrophils by an ethanolic extract of the gum resin exudates of *Boswellia serrata*. *Planta Med* 57:203–207
- Ammon HP, Wahl MA (1991) Pharmacology of *Curcuma longa*. *Planta Med* 57:1–7
- Amos S, Gamaniel K, Akah P, Wambebe C (1998) Anti-inflammatory and muscle relaxant effect of aqueous extract of *Pavetta crassipes* leaves. *Fitoterapia* 69:425–429
- Aqel MB (1991) Relaxant effect of the volatile oil of *Romarinus officinalis* on tracheal smooth muscle. *J Ethnopharmacol* 33:57–62
- Aqel MB, al-Khalil S, Afifi F, al-Eisawi D (1991) Relaxant effects of *Ferula sinaica* root extract on rabbit and guinea pig smooth muscle. *J Ethnopharmacol* 31:373–381
- Arul V, Miyazaki S, Dhananjayan R (2004) Mechanisms of the contractile effect of the alcoholic extract of *Aegle marmelos* Corr on isolated guinea pig ileum and tracheal chain. *Phytomedicine* 11:679–683
- Barns PJ (1989) New concept in the pathogenesis of bronchial hyper-responsiveness and asthma. *J Allergy Clin Immunol* 83:1013–1026
- Barns PJ, Chung KF, Page CP (1988) Inflammatory mediators and asthma. *Pharmacol Rev* 40:49–84
- Baruah CC, Gupta PP, Patnaik GK, Nath A, Kulshreshtha DK, Dhawan BN (1997) Anti-allergic and mast cell stabilizing activity of *Albizia lebbeck*. *Ind Veterinary Med J* 21:127–132
- Bayer T, Breu W, Seligmann O, Wray V, Wagner H (1989) Biologically active thiosulphinates and α -sulphinyl disulphides from *Allium cepa*. *Phytochemistry* 28:2373–2377
- Benancia F, Courreges MC, Coulombie FC (2000) *In vivo* and *in vitro* immunomodulatory activities of *Trichilia glabra* aqueous leaf extracts. *J Ethnopharmacol* 69:199–205
- Benancia F, Courreges MC, Nores MM, Coulombie FC (1995) Immunomodulatory activities of *Cedrela tubiflora* leaf aqueous extracts. *J Ethnopharmacol* 49:133–139
- Bernstein IL (1985) Cromolyn sodium in the treatment of asthma: coming of age in the United States. *J Aller Clin Immunol* 76:381–388
- Campos MG, Toxqui E, Tortoriello J, Oropeza MV, Ponce H, Vargas MH, Montano LM (2001) *Galphimia glauca* organic fraction antagonized LTD (4)-induced contraction in guinea pig airways. *J Ethnopharmacol* 74:7–15
- Chan SC, Chang YS, Wang JP, Chen SC, Kuo SC (1998) Three new flavonoids and anti-allergic, anti-inflammatory constituents from the heartwood of *Dalbergia odorifera*. *Planta Med* 64:153–158
- Channa S, Dar A, Yaqoob M, Anjum S, Sultani Z, Atta-ur-Rahman (2003) Broncho-vasodilatory activity of fractions and pure constituents isolated from *Bacopa monniera*. *J Ethnopharmacol* 86:27–35
- Chao KP, Hua KF, Hsu HY, Su YC, Chang ST (2005) Anti-inflammatory activity of sugiol, a diterpene isolated from *Calocedrus formosana* bark. *Planta Med* 71:300–305
- Chen ZC, Duan XB, Liu KR (1988) The anti allergic activity of osthol extracted from the fruits of *Cnidium monnieri* (L.) Cusson. *Acta Pharmaceutica Sinica* 23:96–99
- Chitravanshi VC, Gupta PP, Kulshreshtha DK, Kar K, Dhawan BN (1990) Anti-allergic activity of *Solanum xanthocarpum*. *Ind J Pharmacol* 22:23–24
- Christie PE, Tagari P, Ford-Hutchinson AW, Black C et al (1992) Urinary leukotriene E₄ after lysine–aspirin inhalation in asthmatic subjects. *Am Rev Respir Dis* 146:1531–1534
- Church MM, Hiroi J (1987) Inhibition of IgE-dependant histamine release from human dispersed lung mast cells by anti-allergic drugs and salbutamol. *Br J Pharmacol* 90:421–429
- Cockcroft DW, Murdock KY (1987) Comparative effects of inhaled salbutamol, sodium chromoglycate and beclomethasone dipropionate on allergen-induced early asthmatic responses, late asthmatic responses and increased bronchial responsiveness to histamine. *J Aller Clin Immunol* 79:734–740
- Cortes SF, Alencar JI, Thomas G, Filho JMB (1995) Spasmolytic action of warifteine, a bisbenzylisoquinoline alkaloid isolated from the root bark of *Cissampelos sympodialis* Eichl. *Phytother Res* 9:579–583
- Cox JSG (1967) Disodium chromoglycate (FPL 670) (Intal): a specific inhibitor of reaginic antibody-antigen mechanisms. *Nature* 216:1328–1329
- Dai Y, Chan YP, Chu LM, Bu PP (2002) Antiallergic and anti-inflammatory properties of the ethanolic extract from *Gleditsia sinensis*. *Biol Pharm Bull* 5:1179–1182
- Dajani BM, Sliman NA, Shubair KS, Hamzeh YS (1981) Bronchospasm caused by intravenous hydrocortisone sodium succinate (Solu-Cortef) in aspirin sensitive asthmatics. *J Allergy Clin Immunol* 68:201–206
- Damre AS, Gokhale AB, Phadke AS, Kulkarni KR, Saraf MN (2003) Studies on the immunomodulatory activity of flavonoidal fraction of *Tephrosia purpurea*. *Fitoterapia* 74:257–261
- Dar A, Channa S (1997) Relaxant effect of ethanolic extract of *Bacopa monniera* on trachea, pulmonary and aorta from rabbit and guinea pig. *Phytother Res* 11:323–325
- Delporte C, Backhouse N, Erazo S, Negrete R, Vidal P, Silva X, Lopez-Perez JL, Feliciano AS, Munoz O (2005) Analgesic–

- antiinflammatory properties of *Proustia pyrifolia*. J Ethnopharmacol 99:119–124
- Dhawan K, Kumar S, Sharma A (2003) Anti-asthmatic activity of the methanol extract of leaves of *Passiflora incarnata*. Phytother Res 17:821–822
- Diamant Z, Timmers MC, Vander Veen H, Friedman BS, De Smet M, Depre M, Hilliard D, Bel EH, Sterk PJ (1995) The effect of MK-0591, a novel 5-lipoxygenase activating protein inhibitor on leukotriene biosynthesis and allergen-induced airway responses in asthmatic subjects *in vivo*. J Aller Clin Immunol 95:42–51
- Dorsch W, Ettl M, Hein G, Scheftner P, Weber J, Bayer T, Wagner H (1987) Anti-asthmatic effects of onions. Inhibition of platelet activating factor induced bronchial obstruction by onion oils. Int Arch Allergy App Immunol 82:535–536
- Drazen JM (1997) Pharmacology of leukotriene receptor antagonists and 5-lipoxygenase inhibitors in the management of asthma. Pharmacotherapy 17:22S–30S
- Dutta NK, Sastry M, Tamhane RG (1968) Pharmacological actions of an alkaloidal fraction isolated from *Saussurea leppa* (Clarke). Curr Sci 37:550–551
- El Gazzar M, El Mezayen R, Nicolls MR, Marecki JC, Dreskin SC (2006) Downregulation of leukotriene biosynthesis by thymoquinone attenuates airway inflammation in a mouse model of allergic asthma. Biochim Biophys Acta 170:1088–1095
- Floreani AA, Rennard SI (1999) The role of cigarette smokes in the pathogenesis of asthma and as a trigger for acute symptoms. Curr Opin Pulm Med 5:38–46
- Geetha VS, Viswanathan S, Kameswaran L (1981) Comparison of total alkaloids of *Tylophora indica* and disodium cromoglycate on mast cell stabilization. Ind J Pharmacol 13:199–201
- Gokhale AB, Dikshit VJ, Damre AS, Kulkarni KR, Saraf MN (2000) Influence of ethanolic extract of *Tephrosia purpurea* Linn. on mast cells and erythrocytes membrane integrity. Ind J Exp Biol 38:837–840
- Goyal BR, Agrawal BB, Goyal RK, Mehta AA (2007) Pharmacological classification of herbal anti-asthmatics. Orient Pharm Exp Med 7:11–25
- Gupta M, Mazumdar UK, Sivakumar T, Vamsi ML, Karki S, Sambathkumar R, Manikandan L (2003) Evaluation of anti-inflammatory activity of chloroform extract of *Bryonia laciniosa* in experimental animal models. Biol Pharm Bull 26:1342–1344
- Gupta SS (1968) Development of anti-histaminic and anti-allergic activity after prolonged administration of a plant saponin from *Clerodendron serratum*. J Pharm Pharmacol 20:801–802
- Gupta SS (1974) Some observations on the anti-asthmatic effect of the saponins of *Gardenia latifolia*. Aspect Aller Appl Immunol 7:198–204
- Gupta SS, Tripathi RM (1973) Effect of chronic treatment of the saponin of *Clerodendron serratum* on disruption of mesenteric mast cells of rats. Aspect Aller Appl Immunol 6:177–188
- Haranath PSRK, Shyamalakumari S (1975) Experimental study on the mode of action of *Tylophora asthmatica* in bronchial asthma. Ind J Med Res 63:661–670
- Harish MS, Mallikarjun N, Badami S (2001) Antihistaminic and mast cell-stabilizing activity of *Striga orobanchioides*. J Ethnopharmacol 76:197–200
- Hashimoto K, Yanagisawa T, Okui Y, Ikeya Y, Maruno M, Fujita T (1994) Studies on anti-allergic components in the roots of *Asiasarum sieboldi*. Planta Med 60:124–127
- Hong SH, Jeong HJ, Kim HM (2003) Inhibitory effects of *Xanthii fructus* extract on mast cell mediated allergic reactions in murine model. J Ethnopharmacol 88:229–234
- Hueza IM, Fonseca ESM, Paulino CA, Haraguchi M, Gorniak SL (2003) Evaluation of immunomodulatory activity of *Ipomoea carnea* on peritoneal cells of rats. J Ethnopharmacol 87:181–186
- Huntley A, Ernst E (2000) Herbal medicines for asthma: a systematic review. Thorax 55:925–29
- Inoue T, Sugimoto Y, Masuda H, Kamei C (2002) Antiallergic effect of flavonoids obtained from *Mentha piperita* L. Biol Pharm Bull 25:256–259
- Ishiguro K, Ueda Y, Iwaoka E, Oku H (2000) Antiallergic and antipruritic effect of *Impatiens textori*. Phytomedicine 7:94–97
- Johri RK, Zutshi U, Kameshwaran L, Atal CK (1985) Effect of quercetin and *Albizia saponins* on rat mast cell. Ind J Physiol Pharmacol 29:43–46
- Kanno M, Shibano T, Takido M, Kitanaka S (1999) Anti-allergic agent from natural sources. 2. structures and leukotriene release-inhibitory effect of toroside B and torosachrysone 8-O-6-malonyl beta gentiobioside from *Cassia torosa* Cav. Chem Pharm Bull 47:915–918
- Kao ES, Wang CJ, Lin WL, Yin YF, Wang CP, Tseng TH (2005) Antiinflammatory potential of flavonoids contents from dried fruits of *Crataegus pinnatifida* *in vitro* and *in vivo*. J Agric Food Chem 53:430–436
- Khalil SA, Aqel M, Afifi F, Eisawi DA (1990) Effect of an aqueous extract of *Ferula ovina* on rabbit and guinea pig smooth muscle. J Ethnopharmacol 30:35–42
- Kim DK, Lee KT, Eun JS, Zee OP, Lim JP, Eum SS, Kim SH, Shin TY (1999) Anti-allergic components from peels of *Citrus unshiu*. Arch Pharm Res 22:642–645
- Kim HM, Lee EH, Hong SH, Song HJ, Shin MK, Kim SH, Shin TY (1998) Effect of *Syzygium aromaticum* extract on immediate hypersensitivity in rats. J Ethnopharmacol 60:125–131
- Kim YC, Lee EH, Lee YM, Kim HK, Song BK, Lee EJ, Kim HM (1997) Effect of the aqueous extract of *Aquillaria agallocha* stem on the immediate hypersensitivity reactions. J Ethnopharmacol 58:31–38
- Kitanaka S, Nakayama T, Shibano T, Ohkoshi E, Takido M (1998) Anti-allergic agent from natural sources, structures and inhibitory effect of histamine release of naphthopyrone glycosides from seeds of *Cassia obtusifolia* L. Chem Pharm Bull 46:1650–1652
- Kostalova D, Bukovsky M, Koscova H, Kardosova A (2001) Anticomplement activity of *Mahonia aquigolium* bisbenzylisoquinoline alkaloids and berberine extract. Ceska Slov Farm 50:286–289
- Kou J, Sun Y, Lin Y, Cheng Z, Zheng W, Yu B, Xu Q (2005) Antiinflammatory activities of aqueous extract from radix *Ophiopogon japonicus* and its two constituents. Biol Pharm Bull 28:1234–1238
- Kubo M, Matsuda H, Tomohiro N, Yoshikawa M (1997) Studies on *Alismatis rhizome*; Anti-allergic effects of methanol extract and six terpene components from *Alismatis rhizoma* (Dried rhizome of *Alisma orientale*). Biol Pharm Bull 20:511–516
- Kumar DA, Ramu P (2002) Effect of methanolic extract of *Benincasa hispida* against histamine and acetylcholine induced bronchospasm in guinea pigs. Ind J Pharmacol 34:365–366
- Kumar Suresh (1979) Scientific Appraisal of *Adhatoda vasica* _Nees (Vasaka) J NIMA XXIII: 257–261
- Kumar VL, Basu N (1994) Anti-inflammatory activity of the latex of *Calotropis procera*. J Ethnopharmacol 44:123–125
- Lee J, Jung E, Park J, Jung K, Lee S, Hong S, Park J, Park E, Kim J, Park S, Park D (2005) Antiinflammatory effects of magnolol and honokiol are mediated through inhibition of the downstream pathway of MEKK-1 in NR-kB activation signaling. Planta Med: 71:338–343
- Makare N, Bodhankar S, Rangari V (2001) Immunomodulatory activity of alcoholic extract of *Mangifera indica* L. in mice. J Ethnopharmacol 78:133–137
- Mali RG, Mahajan SG, Mehta AA (2008) Studies on Bronchodilatory effect of *Lepidium sativum* against allergen induced bronchospasm in Guinea pigs. Pharmacog Mag 4:189–192

- Manez S, Alcaraz MJ, Paya M, Rios JL, Hancke JL (1990) Selected extracts from medicinal plants as anti-inflammatory agents. *Planta Med* 56:656
- Manosroi A, Saraphanchotiwithaya A, Manosroi J (2005) *In vivo* immunomodulating activity of wood extracts from *Clausena excavate* Burm.F. *J Ethnopharmacol* 102:5–9
- Marone G, Columbo M, Triggiani M, Cirillo R, Genovese A, Formisano S (1987) Inhibition of IgE mediated release of histamine and peptide leukotriene from human basophils and mast cells by forskolin. *Biochem Pharmacol* 36:13–20
- Matsuda H, Shimoda H, Yamahara J, Yoshikawa M (1999) Effect of phyllodulcin, hydrangenol, and their 8-O-glucosides, and thunberginol A and F from *Hydrangea macrophylla* var. thunbergii on passive cutaneous anaphylaxis reaction in rats. *Biol Pharm Bull* 22:870–872
- Matsuda H, Tomohiro N, Ido Y, Kubo M (2002) Anti-allergic effects of *Cnidii monnieri* (dried fruits of *Cnidium monnieri*) and its major component, osthol. *Biol Pharm Bull* 25:809–812
- Meister A, Bernhard G, Chrisoffel V, Buschauer A (1999) Antispasmodic activity of *Thymus vulgaris* extract on isolated g.pig trachea: discrimination between drug and ethanol effects. *Planta Med* 65:512–516
- Mengi SA, Deshpande SG (1999) Anti-inflammatory activity of *Butea frondosa* leaves. *Fitoterapia* 70:521–522
- Moran A, Carron R, Martin ML, San Roman L (1989) Anti-asthmatic activity of *Artemisia caerulea* subsp. gallica. *Planta Med* 55:351–353
- Mukherjee PK, Saha K, Bhattacharya S, Giri SN, Pal M, Saha BP (1997) Studies on antitussive activity of *Drymaria cordata* Willd. *J Ethnopharmacol* 56:77–80
- Muller A, Antus S, Bittinger M, Dorsch W, Kaas A, Kreher B et al (1993) Chemistry and pharmacology of the antiasthmatic plants *Galphimia glauca*, *Adhatoda vasica* and *Picrorrhiza kurroa*. *Planta Med* 59(A5):86
- Mungantiwar AA, Nair AM, Shinde UA, Dikshit VJ, Saraf MN, Thakur VS, Sainis KB (1999) Studies on the immunomodulatory effects of *Boerhaavia diffusa* alkaloidal fraction. *J Ethnopharmacol* 65:125–131
- Nair AM, Saraf MN (1995) Inhibition of antigen and compound 48/80 induced contractions of guinea pig trachea by the ethanolic extract of the leaves of *Vitex negundo* Linn. *Ind J Pharmacol* 27:230–233
- Nair AM, Tamhankar CP, Saraf MN (1994) Studies on the mast cell stabilizing activity of *Vitex negundo* Linn. *Ind Drugs* 32:277–282
- National Institute of Health (1997) Expert Panel Report 2. Guidelines for the diagnosis and Management of asthma. NIH Publication. No.97-4051
- Nayampalli S, Desai NK, Ainapure SS (1986) Anti-allergic properties of *Tinospora cordifolia* in animal models. *Ind J Pharmacol* 18:250–252
- Neszmelyi A, Kreher B, Muller A, Dorsch W, Wagner H (1993) Tetragalloylquinic acid, the major Antiasthmatic principle of *Galphimia glauca*. *Planta Med* 59:164–167
- Okpo SO, Adeyemi OO (2002) The anti-allergic effects of *Crinum glaucum* aqueous extract. *Phytomedicine* 9:438–441
- Oli RG, Manikandan L, Swarna FB, Manikandan P, Khosa RL (2005) Evaluation of antiinflammatory potential of *Indigofera tinctoria* extract in rats. *Ind J Nat Prod* 21:12–15
- Owoyele VB, Oloriegebe YY, Balogun EA, Soladoye AO (2005) Analgesic and antiinflammatory properties of *Nelsonia canescens* leaf extract. *J Ethnopharmacol* 99:153–156
- Palanichamy S, Amala Bhaskar E, Nagarajan S (1991) Effect of *Cassia alata* leaf extract on mast cell stabilization. *Ind J Pharmacol* 23:189–191
- Paliwa JK, Dwiwedi AK, Singh S (2000) Pharmacokinetics and in-situ absorption studies of a new anti-allergic compound 73/602 in rats. *Int J Pharm* 197:213–220
- Pavia D, Batement JRM, Clarke SW (1980) Deposition and clearance of inhaled particles. *Bull Eur Physiopath Resp* 16:335–366
- Persson CG, Draco AB (1988) Xanthine as airway anti-inflammatory drugs. *J Allergy Clin Immunol* 81:615–617
- Puglisi L, Salvadori S, Gabrielli G, Pasargiklian R (1988) Pharmacology of natural compounds. Smooth muscle relaxant activity induced by a Ginkgo biloba L. extract on guinea-pig trachea. *Pharmacol Res Comm* 20:573–589
- Qin F, Sun HX (2005) Immunosuppressive activity of pollen *Typhae* ethanol extract on the immune responses in mice. *J Ethnopharmacol* 102:424–429
- Rao YK, Fang SH, Tzeng YM (2005) Inhibitory effects of the flavonoids isplated from *Waltheria indica* on the production of NO, TNF- α and IL-2 in activated macrophages. *Biol Pharm Bull* 28:912–915
- Rasool M, Varalakshmi P (2006) Immunomodulatory role of *Withania somnifera* root powder on experimental induced inflammation: An *in vivo* and *in vitro* study. *Vascul Pharmacol* 44:406–410
- Rezaeipoor R, Saeidnia S, Kamalinejad M (2000) The effects of *Plantago ovata* on humoral immune responses in experimental animals. *J Ethnopharmacol* 72: 283–286
- Samiulla DS, Prashanth D, Amit A (2001) Mast-cell stabilizing activity of *Bacopa monnieri*. *Fitoterapia* 72:284–285
- Saraf MN, Patwardhan BK (1988a) Pharmacological studies on *Sarcostemma brevistigma*. Part I Anti-allergic activity. *Ind Drugs* 26:49–53
- Saraf MN, Patwardhan BK (1988b) Pharmacological studies on *Sarcostemma brevistigma*. Part II Bronchodilator activity. *Ind Drugs* 26:54–57
- Sayah ME, Filho VC, Yunes RA, Pinheiro TR, Calixto JB (1998) Action of polygodial, a sesquiterpene isolated from *Drymis winteri* in the guinea-pig ileum an trachea *in vitro*. *Eur J Pharmacol* 344:215–221
- Sen P (1993) Therapeutic potential of Tulsi (*Ocimum sanctum*) from experience to fact. *Drug Views* 1:15–18
- Shin TY, Jeong HJ, Kim DK, Kim SH, Lee JK, Chae BS, Kim JH, Kang HW, Lee CM (2001a) Inhibitory action of water-soluble fraction of *Terminalia chebula* on systemic and local anaphylaxis. *J Ethnopharmacol* 74:133–140
- Shin TY, Kim DK, Chae BS, Lee EJ (2001b) Antiallergic action of *Magnolia officinalis* on immediate hypersensitivity reaction. *Arch Pharm Res* 24:249–255
- Shin TY, Kim SH, Lim JP, Suh ES, Jeong HJ, Kim BD, Park EJ, Hwang WJ, Rye DG, Baek SH, An NH, Kim HM (2000) Effect of *Vitex rotundifolia* on immediate-type allergic reaction. *J Ethnopharmacol* 72:443–450
- Shin TY, Kim SH, Suk K, Ha JH, Kim I, Lee MG, Jun CD, Kim SY, Lim JP, Eun JS, Shin HY, Kim HM (2005) Anti-allergic effects of *Lycopus lucidus* on mast cellmediated allergy model. *Toxicol Appl Pharmacol* 209:255–262
- Shin TY, Park JH, Kim HM (1999) Effect of *Cryptotympana atrata* extract on compound 48/80 induced anaphylactic reactions. *J Ethnopharmacol* 66:319–325
- Shinde UA, Phadke AS, Kulkarni KR, Nair AM, Mungantiwar AA, Dikshit VJ, Saraf MN (1999) Mast cell stabilizing and lipooxygenase inhibiting activity of *Cedrus deodara* (Roxb.) wood oil. *Ind J Exp Biol* 37:258–261
- Singh RK, Acharya SB, Bhattacharya SK (2000) Pharmacological activity of *Elaeocarpus sphaericus*. *Phytother Res* 14:36–39
- Singh S, Agrawal SS (1990) Broncho-relaxant activity of *Belamcanda chinensis*. *Ind J Pharmacol* 22:107–109
- Singh S, Agrawal SS (1991) Anti asthmatic and anti-inflammatory activity of *Ocimum sanctum*. *Int J Pharmacog* 29:306–310
- Smit HF, Kroes BH, van den Berg AJJ, van der Wal D, van den Worm E, Beukelman CJ, van Dijk H, Labadie RP (2000) Immunomodulatory

- and anti-inflammatory activity of *Picrorrhiza scrophulariiflora*. J Ethnopharmacol 73:101–109
- Soares de Moura R, Costa SS, Jansen JM, Silva CA, Lopes CS, Bernardo-Filho M, Nascimento da Silva V, Criddle DN, Portela BN, Rubenich LM, Araujo RG, Carvalho LC (2002) Bronchodilator activity of *Mikania glomerata* Sprengel on human bronchi and guinea-pig trachea. J Pharm Pharmacol 54:249–256
- Srivastava S, Gupta PP, Prasad R, Dixit KS, Palit G, Ali B, Mishra G, Saxena RC (1999) Evaluation of anti-allergic activity (Type I hypersensitivity) of *Inula racemosa* in rats. Ind J Physiol Pharmacol 43:235–241
- Stuppner H, Dorsch W, Wagner H, Gropp M, Kepler P (1991) Antiasthmatic effects of *Picrorrhiza kurroa*: inhibition of allergen and PAF induced bronchial obstruction in g.pigs by Androsin, Apocynine and structurally related compounds. Planta Med 57: A62
- Subramoniam A, Evans DA, Valsaraj R, Rajasekharan S, Pushpangadan P (1999) Inhibition of antigen-induced degranulation of sensitized mast cells by *Trichopus zeylanicus* in mice and rats. J Ethnopharmacol 68:137–143
- Suresh Kumar, R.N. Dwivedi and G. N. Chaturvedi, (1981): Scientific Appraisal of Albizzialebbeck-Benth (Shirisha), *J.NIMA*, XXIII, 311–316
- Suzuki M, Yoshino K, Yamamoto MM, Miyase T, Sano M (2000) Inhibitory effect of Tea catechins and o-methylated derivatives of (–)—epigallocatechin –3-O-gallate on mouse type IV allergy. J Agri Food Chem 48:5649–5653
- Tamaoki J, Kondo M, Sakai N (1997) Leukotriene antagonist prevents exacerbation of asthma during reduction of high dose inhaled corticosteroids. Am J Respir Crit Care Med 155:1235–1240
- Thomas G, Araujo CC, Agra MF, Diniz M (1995) Preliminary studies on the hydroalcoholic extract of the root of *Cissampelos sympodialis* Eichl in guinea pig tracheal strips and bronchoalveolar leucocytes. Phytother Res 9:473–477
- Thomas G, Araujo CC, Duarte JC, De souza DP (1997) Bronchodilator activity of an aqueous fraction of an ethanol extract of the leaves of *Cissampelos sympodialis* Eichl. in the guinea pig. Phytomedicine 4:233–238
- Tiwari U, Rastogi B, Thakur S, Jain S, Jain NK (2003) Studies on the immunomodulatory effects of *Cleome viscosa*. Indian J Pharm Sci 66:171–176
- Touvy C, Eienne A, Braquet P (1985) Inhibition of antigen induced lung anaphylaxis in the guinea pig by BN 52021 a new specific PAF-acether receptor antagonist isolated from *Ginkgo biloba*. Agents Actions 17:371–372
- Tripathi RM, Das PK (1977) Studies on anti-asthmatic and anti-anaphylactic activity of *Albizzia lebbeck*. Ind J Pharmacol 9:189–194
- Tripathi RM, Sen PC, Das PK (1979) Studies on the mechanism of action of *Albizzia lebbeck*, an Indian indigenous drug used in the treatment of atopic allergy. J Ethnopharmacol 1:385–386
- Udapa AL, Udapa SL, Guruswamy MN (1991) The possible site of anti-asthmatic action of *Tylophora asthmatica* on pituitary-adrenal axis in albino rats. Planta Med 57:409–413
- Ueda Y, Oku H, Iinuma M, Ishiguro K (2003) Effect on blood pressure decrease in response to PAF of *Impatiens textori*. Biol Pharm Bull 26:1505–1507
- Vázquez B, Avila G, Segura D, Escalante B (1996) Anti-inflammatory activity of extracts from *Aloe vera* gel. J Ethnopharmacol 55:69–75
- Vieira-de-Abreu A, Amendoeira FC, Gomes GS, Zanon C, Chedier LM, Figueiredo MR, Kaplan MA, Frutuosa VS, Castro-Faria-Neto HC, Weller PF, Bandeira-Melo C, Bozza PT (2005) Anti-allergic properties of the bromeliaceae *Nidularium procerum*: inhibition of eosinophil activation and influx. Int Immunopharmacol 5:1966–1974
- Wang N, Yao X, Ishii R, Kitanaka S (2001) Antiallergic agents from natural sources: structures and inhibitory effects on nitric oxide production and histamine release of five novel polyacetylene glucosides from *Bidens parviflora* Willd. Chem Pharm Bull 49:938–942
- Weinberger M (1984) The pharmacology and therapeutic use of theophylline. J Allergy Clin Immunol 73:525–540
- Wenzel SE, Trudeau JB, Kaminsky DA, Cohn J, Martin RJ, Westcott JY (1995) Effect of 5-lipoxygenase inhibition on bronchoconstriction and airway inflammation in nocturnal asthma. Am J Respir Crit Care Med 152:897–905
- Wu JB, Chun YT, Ebizuka Y, Sankawa V (1985) Biologically active constituents of *Centipeda minima*: isolation of a new sesquiterpene lactones. Chem Pharm Bull 33:4091–4094
- Yang T, Jia M, Meng J, Wu H, Mei Q (2006) Immunomodulatory activity of polysaccharides isolated from *Angelica sinensis*. Int J Biol Macromol 39:179–184
- Yi JM, Hong SH, Kim JH, Kim HK, Song HJ, Kim HM (2002) Effect of *Acanthopanax senticosus* stem on mast cell-dependant anaphylaxis. J Ethnopharmacol 79:347–352